

R. Review Materials

Contents

R1 Mathematical Formulas and Identities	1
R1.1 Finite and Infinite Sums of Numbers ¹	1
R1.2 Power Series	2
R1.3 Factorial	3
R1.4 Permutations and Combinations	3
R1.5 Polynomial Factors and Products	3
R1.6 Roots of Quadratic Equation	4
R1.7 Euler's Formula	4
R1.8 Trigonometric Functions and Formulas	5
R1.9 Newton-Raphson Method: Finding a root of a polynomial equation	6
R1.10 Hölder's Inequality and Cauchy-Schwartz's Inequality	7
R2 Useful Functions	9
R3 Commonly Used Differentials and Integrals	11
R3.1 Differentials	11
R3.2 Integrals	11
R3.3 l'Hôpital's Rule	12
R3.4 Examples	12
R4 Complex Numbers	15
R4.1 Definition	15
R4.2 Complex Arithmetic	16
R5 Complex Variables	18
R5.1 Function of a Complex Variable	18
R5.2 Analytic Function of a Complex Variable	18
R5.3 Analytic Continuation	19
R5.4 Cauchy's Integral Formula	19
R5.5 Cauchy's Residue Theorem	19

R6 Continuous-Time Signals	20
R6.1 Energy and Power	20
R6.2 Continuous-Time Sinusoidal and Exponential Signals	20
R6.2.1 Definition	20
R6.2.2 Properties	21
R6.3 Continuous-Time Eigenfunction	22
R6.4 Continuous-Time Fourier Series	23
R6.4.1 Definition	23
R6.4.2 Dirichlet Conditions	25
R6.5 Continuous-Time Fourier Transform	26
R7 Discrete Fourier Series	27
R8 Matrix Algebra	29
R8.1 Definition	29
R8.2 Transpose	29
R8.3 Toeplitz Matrix	30
R8.4 Circulant Matrix	30
R8.5 Determinant	30
R8.6 Minor and Cofactor	32
R8.7 Inverse of a Matrix	33
R8.8 Unitary Matrix and Orthogonal Matrix	33
R8.9 Cramer's Rule	34

R1 Mathematical Formulas and Identities

R1.1 Finite and Infinite Sums of Numbers¹

$$\sum_{k=1}^n k = \frac{n(n+1)}{2}, \quad (\text{R1.1})$$

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}, \quad (\text{R1.2})$$

$$\sum_{k=1}^n k^3 = \frac{n^2(n+1)^2}{4}, \quad (\text{R1.3})$$

$$\sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}, \quad (\text{R1.4})$$

$$\sum_{k=1}^{\infty} \frac{1}{k^4} = \frac{\pi^4}{90}, \quad (\text{R1.5})$$

where n is a positive integer.

Note: The series

$$\sum_{k=1}^{\infty} \frac{1}{k} \quad (\text{R1.6})$$

does not converge.

$$\sum_{k=0}^{n-1} d^k = \frac{1-d^n}{1-d}, \quad (\text{R1.7})$$

with d arbitrary integer and $|d| \neq \{1, 0\}$.

$$\sum_{k=0}^{\infty} d^k = \frac{1}{1-d}, \quad (\text{R1.8})$$

with $0 < |d| < 1$.

¹For test of the convergence of infinite sums, a recommended reading is *Table of Integrals, Series, and Products*, I.S. Gradshteyn and I.M. Ryzhik, ©2000, Academic Press.

R1.2 Power Series

Binomial Series:

$$(x + y)^n = x^n + \binom{n}{1}x^{n-1}y + \binom{n}{2}x^{n-2}y^2 + \binom{n}{3}x^{n-3}y^3 + \cdots + \binom{n}{n-1}xy^{n-1} + y^n, \quad (\text{R1.9})$$

with n a positive integer.

Taylor Series:

If $f(x)$ is an arbitrarily differentiable function, then it can be expressed in the form

$$f(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2 + \cdots + \frac{f^{(n)}(a)}{n!}(x - a)^n + \cdots, \quad (\text{R1.10})$$

where $f'(a) = \left. \frac{df(x)}{dx} \right|_{x=a}$, $f''(a) = \left. \frac{d^2f(x)}{dx^2} \right|_{x=a}$, and $f^{(n)}(a) = \left. \frac{d^n f(x)}{dx^n} \right|_{x=a}$.

Exponential Series:

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots, \quad (\text{R1.11})$$

$$a^x = 1 + x \log_e a + \frac{(x \log_e a)^2}{2!} + \frac{(x \log_e a)^3}{3!} + \cdots. \quad (\text{R1.12})$$

Logarithmic Series:

$$\log_e(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \cdots, \quad (\text{Region of Convergence : } -1 < x < 1). \quad (\text{R1.13})$$

Trigonometric Series:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots, \quad (\text{R1.14})$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots, \quad (\text{R1.15})$$

$$\tan x = x + \frac{x^3}{3} + \frac{2x^5}{15} + \frac{17x^7}{315} + \frac{62x^9}{2835} + \cdots, \quad (\text{Region of Convergence : } x^2 < \frac{\pi^2}{4}), \quad (\text{R1.16})$$

$$\cot x = \frac{1}{x} - \frac{x}{3} - \frac{x^3}{45} - \frac{2x^5}{945} + \cdots, \quad (\text{Region of Convergence : } 0 < |x| < \pi). \quad (\text{R1.17})$$

R1.3 Factorial

$$n! = n(n-1)(n-2)\cdots 2 \cdot 1, \quad \text{with } n \text{ a nonnegative integer}, \quad (\text{R1.18})$$

$$0! = \Gamma(0+1) = 1. \quad (\text{R1.19})$$

R1.4 Permutations and Combinations

The number of permutations S of n things taken k at a time, with n and k positive integers, is given by

$$S = \frac{n!}{(n-k)!}. \quad (\text{R1.20})$$

The number of combinations S of n things taken k at a time, with n and k positive integers, is given by

$$S = \binom{n}{k} = \frac{n!}{k!(n-k)!}. \quad (\text{R1.21})$$

R1.5 Polynomial Factors and Products

$$x^n - y^n = (x-y)(x^{n-1} + x^{n-2}y + \cdots + y^{n-1}), \quad (\text{R1.22})$$

with n a positive integer.

$$x^n + y^n = (x + y)(x^{n-1} - x^{n-2}y + x^{n-3}y^2 - \cdots + y^{n-1}), \quad (\text{R1.23})$$

with n a positive and odd integer.

$$\begin{aligned} \prod_{i=1}^N (x + \xi_i) &= (x + \xi_1)(x + \xi_2) \cdots (x + \xi_N) \\ &= \alpha_0 + \alpha_1 x + \alpha_2 x^2 + \cdots + \alpha_{N-1} x^{N-1} + \alpha_N x^N \end{aligned} \quad (\text{R1.24})$$

where

$$\begin{aligned} \alpha_0 &= \prod_{i=1}^N \xi_i, & \alpha_1 &= \sum_{i=1}^N \frac{\alpha_0}{\xi_i}, & \alpha_2 &= \sum_{\substack{i \neq j \\ i, j=1}}^N \frac{\alpha_0}{\xi_i \xi_j}, & \cdots, \\ \alpha_{N-1} &= \xi_1 + \xi_2 + \xi_3 + \cdots + \xi_N, & \alpha_N &= 1. \end{aligned}$$

R1.6 Roots of Quadratic Equation

The roots x_1, x_2 of the quadratic equation

$$ax^2 + bx + c = 0,$$

with a, b , and c real numbers, are given by

$$x_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}, \quad (\text{R1.25})$$

$$x_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}. \quad (\text{R1.26})$$

Note:

$$x_1 + x_2 = \frac{-b}{a}, \quad (\text{R1.27})$$

$$x_1 x_2 = \frac{c}{a}. \quad (\text{R1.28})$$

R1.7 Euler's Formula

$$e^{j\theta} = \cos \theta + j \sin \theta, \quad (\text{R1.29})$$

with θ a real number.

R1.8 Trigonometric Functions and Formulas

$$\sin \theta = \frac{1}{2j}(e^{j\theta} - e^{-j\theta}), \quad (\text{R1.30})$$

$$\cos \theta = \frac{1}{2}(e^{j\theta} + e^{-j\theta}), \quad (\text{R1.31})$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{(e^{j\theta} - e^{-j\theta})}{j(e^{j\theta} + e^{-j\theta})}, \quad (\text{R1.32})$$

$$\cot \theta = \frac{1}{\tan \theta} = \frac{j(e^{j\theta} + e^{-j\theta})}{(e^{j\theta} - e^{-j\theta})}, \quad (\text{R1.33})$$

$$\csc \theta = \frac{1}{\sin \theta} = \frac{2j}{e^{j\theta} - e^{-j\theta}}, \quad (\text{R1.34})$$

$$\sec \theta = \frac{1}{\cos \theta} = \frac{2}{e^{j\theta} + e^{-j\theta}}, \quad (\text{R1.35})$$

$$\sin \theta = \cos\left(\frac{\pi}{2} - \theta\right) = \sin(\pi - \theta), \quad (\text{R1.36})$$

$$\cos \theta = \sin\left(\frac{\pi}{2} - \theta\right) = -\cos(\pi - \theta), \quad (\text{R1.37})$$

$$\tan \theta = \cot\left(\frac{\pi}{2} - \theta\right) = -\tan(\pi - \theta), \quad (\text{R1.38})$$

$$\sinh \theta = \frac{1}{2}(e^{\theta} - e^{-\theta}), \quad (\text{R1.39})$$

$$\cosh \theta = \frac{1}{2}(e^{\theta} + e^{-\theta}), \quad (\text{R1.40})$$

$$\tanh \theta = \frac{\sinh \theta}{\cosh \theta} = \frac{(e^{\theta} - e^{-\theta})}{(e^{\theta} + e^{-\theta})}, \quad (\text{R1.41})$$

with θ a real number.

$$\sin(\theta_1 \pm \theta_2) = \sin \theta_1 \cos \theta_2 \pm \cos \theta_1 \sin \theta_2, \quad (\text{R1.42})$$

$$\cos(\theta_1 \pm \theta_2) = \cos \theta_1 \cos \theta_2 \mp \sin \theta_1 \sin \theta_2, \quad (\text{R1.43})$$

$$\sin^2 \theta_1 - \sin^2 \theta_2 = \sin(\theta_1 + \theta_2) \cdot \sin(\theta_1 - \theta_2), \quad (\text{R1.44})$$

$$\cos^2 \theta_1 - \cos^2 \theta_2 = -\sin(\theta_1 + \theta_2) \cdot \sin(\theta_1 - \theta_2), \quad (\text{R1.45})$$

$$\cos^2 \theta_1 - \sin^2 \theta_2 = \cos(\theta_1 + \theta_2) \cdot \cos(\theta_1 - \theta_2), \quad (\text{R1.46})$$

$$\cos^2 \theta_1 + \sin^2 \theta_2 = 1 \quad (\text{R1.47})$$

$$\sin \theta_1 \pm \sin \theta_2 = 2 \sin\left(\frac{\theta_1 \pm \theta_2}{2}\right) \cdot \cos(\theta_1 \mp \theta_2), \quad (\text{R1.48})$$

$$\cos \theta_1 + \cos \theta_2 = 2 \cos\left(\frac{\theta_1 + \theta_2}{2}\right) \cdot \cos\left(\frac{\theta_1 - \theta_2}{2}\right), \quad (\text{R1.49})$$

$$\cos \theta_1 - \cos \theta_2 = -2 \sin\left(\frac{\theta_1 + \theta_2}{2}\right) \cdot \sin\left(\frac{\theta_1 - \theta_2}{2}\right), \quad (\text{R1.50})$$

$$\sin 2\theta = 2 \sin \theta \cos \theta, \quad (\text{R1.51})$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta, \quad (\text{R1.52})$$

$$\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta, \quad (\text{R1.53})$$

$$\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta. \quad (\text{R1.54})$$

with θ , θ_1 , and θ_2 real numbers.

R1.9 Newton-Raphson Method: Finding a root of a polynomial equation

The Newton-Raphson method is a numerical technique to determine approximately the root of the equation $f(x) = 0$. The procedure starts from an initial guess of the root $x = x_1$. Then using the recurrence relation

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}, \quad n = 1, 2, \dots,$$

where

$$f'(x_n) = \left. \frac{df(x)}{dx} \right|_{x=x_n},$$

the successive approximations x_{n+1} , $n \geq 1$, beginning with $n = 1$, can be found. The approximation is assumed to converge when the difference between x_{n+1} and x_n is below a prescribed small number, typically 10^{-6} .

The Newton-Raphson method converges fast to the actual root if the initial guess of the root is close to the actual root. However, there are three main drawbacks: (1) The method fails when $f'(x_n) = 0$, (2) The method does not always converge, and (3) The method may converge to a root different from that expected if the initial guess x_1 is far from the actual root.

Example R1.1. In this example, we would like to show how the Newton-Raphson method is used to find the root of $f(x) = x^3 - 3x^2 + x - 1 = 0$. Assume the numerical resolution required is 14 decimal digits.

We start with an initial guess of the root $x_1 = 2.5$:

$$\begin{aligned} x_1 &= 2.5, \\ x_2 &= x_1 - \frac{f(x_1)}{f'(x_1)} = 2.84210526315789, \\ x_3 &= x_2 - \frac{f(x_2)}{f'(x_2)} = 2.77282691999216, \\ x_4 &= x_3 - \frac{f(x_3)}{f'(x_3)} = 2.76930129255045, \\ x_5 &= x_4 - \frac{f(x_4)}{f'(x_4)} = 2.76929235429601, \\ x_6 &= x_5 - \frac{f(x_5)}{f'(x_5)} = 2.76929235423863, \\ x_7 &= x_6 - \frac{f(x_6)}{f'(x_6)} = 2.76929235423863. \end{aligned}$$

The recurrence process stops as $|x_7 - x_6| \leq 10^{-15}$. Hence $x = x_7$ is a root of $f(x)$.

R1.10 Hölder's Inequality and Cauchy-Schwartz's Inequality

The Hölder's inequality for integrals is given by

$$\left| \int_{a_0}^{a_1} f(x)g(x) dx \right| \leq \left(\int_{a_0}^{a_1} |f(x)|^p dx \right)^{1/p} \left(\int_{a_0}^{a_1} |g(x)|^q dx \right)^{1/q}, \quad (\text{R1.55})$$

where

$$\frac{1}{p} + \frac{1}{q} = 1.$$

The equality holds when

$$f(x) = kg(x)^{p-1}, \text{ with } k \text{ any constant.}$$

If $p = q = 2$, the inequality becomes Schwartz's inequality

$$\left| \int_{a_0}^{a_1} f(x)g(x) dx \right| \leq \left(\int_{a_0}^{a_1} |f(x)|^2 dx \right)^{1/2} \left(\int_{a_0}^{a_1} |g(x)|^2 dx \right)^{1/2}. \quad (\text{R1.56})$$

The equality holds when

$$f(x) = kg(x), \text{ with } k \text{ any constant.}$$

The Hölder's inequality for sums is given by

$$\left| \sum_{i=1}^N x_i y_i \right| \leq \left(\sum_{i=1}^N |x_i|^p \right)^{1/p} \left(\sum_{i=1}^N |y_i|^q \right)^{1/q}, \quad (\text{R1.57})$$

where

$$\frac{1}{p} + \frac{1}{q} = 1.$$

The equality holds when

$$y_i = kx_i^{p-1}, \text{ with } k \text{ any constant.}$$

If $p = q = 2$, the inequality becomes Cauchy's inequality

$$\left| \sum_{i=1}^N x_i y_i \right| \leq \left(\sum_{i=1}^N |x_i|^2 \right)^{1/2} \left(\sum_{i=1}^N |y_i|^2 \right)^{1/2}, \quad (\text{R1.58})$$

The equality holds when

$$y_i = kx_i, \text{ with } k \text{ any constant.}$$

R2 Useful Functions

1. *rect* function

$$\text{rect}(x) = \begin{cases} 1, & |x| < \frac{1}{2} \\ 0, & |x| > \frac{1}{2} \end{cases}.$$

2. *sinc* function

$$\text{sinc}(x) = \frac{\sin \pi x}{\pi x}.$$

3. *signum* function

$$\text{sgn}(x) = \begin{cases} +1, & x > 0 \\ 0, & x = 0 \\ -1, & x < 0 \end{cases}.$$

4. *ceiling* function rounds the input x towards the closest integer larger than or equal to x and is denoted as $\lceil x \rceil$.

For example, $\lceil 3.2 \rceil = \lceil 3.8 \rceil = 4$ and $\lceil -3.2 \rceil = \lceil -3.9 \rceil = -3$.

5. *floor* function rounds the input x towards the closest integer less than or equal to x and is denoted as $\lfloor x \rfloor$.

For example, $\lfloor 3.2 \rfloor = \lfloor 3.8 \rfloor = 3$ and $\lfloor -3.2 \rfloor = \lfloor -3.9 \rfloor = -4$.

6. *median* of a set of real numbers $\{x_1, x_2, \dots, x_N\}$ is obtained by rank ordering the numbers in the set and choosing the middle number in the ordered set.

For example, the median of $\{7, 13, 1, 6, 3\}$ is 6 and the median of $\{7, 13, 1, 6, 3, 9\}$ is $(6 + 7)/2 = 6.5$.

7. *Dirac delta* function $\delta(\tau)$ is a function of τ with infinite height, zero width, and unit area. It is the limiting form of a unit area pulse function

$$p_{\Delta}(\tau) = \begin{cases} \frac{1}{2\Delta}, & -\Delta < \tau \leq \Delta \\ 0, & \text{elsewhere} \end{cases}$$

as Δ goes to 0, i.e.,

$$\lim_{\Delta \rightarrow 0} \int_{-\infty}^{\infty} p_{\Delta}(\tau) d\omega = \int_{-\infty}^{\infty} \delta(\tau) d\tau = 1. \quad (\text{R2.1})$$

Equation(R2.1) also holds even when we reverse the direction of axis τ and shift $\delta(-\tau)$ by an amount t , i.e.,

$$\int_{-\infty}^{\infty} \delta(t - \tau) d\tau = 1. \quad (\text{R2.2})$$

Because of the above properties, we have

$$\int_{-\infty}^{\infty} x(\tau)\delta(t - \tau) d\tau = x(\tau)|_{\tau=t} = x(t). \quad (\text{R2.3})$$

Equation(R2.3) holds for any value of t , and it is referred as the *sifting property* of the Dirac delta function.

8. The *modulo* operation of integer X over integer N is the residue of X divided by N :

$$\langle X \rangle_N = X - kN, k = \lfloor X/N \rfloor.$$

It can be verified that the modulo operation is *linear*. When negative numbers are used, $\langle X \rangle_N$ has the same sign as N . For example, $\langle 67 \rangle_{13} = 67 - 5 \cdot 13 = 2$, $\langle 67 \rangle_{-13} = 67 - (-13) \cdot (-6) = -11$ and $\langle -67 \rangle_{13} = -67 - 13 \cdot (-6) = 11$.

The statement “ X is congruent to Y , modulo N ” means that

$$\langle X \rangle_N = \langle Y \rangle_N.$$

The notation

$$\langle X^{-1} \rangle_N$$

denotes the multiplicative inverse of X evaluated modulo N , i.e., if $\langle X^{-1} \rangle_N = \alpha$, then $\langle X\alpha \rangle_N = 1$. For example, $\langle 3^{-1} \rangle_4 = 3$ because $\langle 3 \cdot 3 \rangle_4 = 1$, and $\langle 8^{-1} \rangle_5 = 2$ because $\langle 8 \cdot 2 \rangle_5 = 1$.

In the case of polynomial, the operation $a(z) \bmod b(z)$ is the residue $r(z)$ after the polynomial division $a(z)/b(z)$. For example, if $a(z) = 4z^{-3} + 2z^{-2} + 5z^{-1} + 1$ and $b(z) = z^{-2} + 3z^{-1} + 4$ then the residue after the division

$$\frac{a(z)}{b(z)} = 4z^{-1} - 10 + \frac{19z^{-1} + 41}{z^{-2} + 3z^{-1} + 4}$$

is $19z^{-1} + 41$. Therefore, $a(z) \bmod b(z) = r(z) = 19z^{-1} + 41$.

R3 Commonly Used Differentials and Integrals

R3.1 Differentials

$$d(uv) = u dv + v du \quad (\text{R3.1})$$

$$d\left(\frac{u}{v}\right) = \frac{v du - u dv}{v^2} \quad (\text{R3.2})$$

$$d(u^n) = n u^{n-1} du \quad (\text{R3.3})$$

$$d e^u = e^u du \quad (\text{R3.4})$$

$$d a^u = (a^u \log_e a) du \quad (\text{R3.5})$$

$$d(\log_e u) = u^{-1} du \quad (\text{R3.6})$$

$$d \sin u = \cos u du \quad (\text{R3.7})$$

$$d \cos u = -\sin u du. \quad (\text{R3.8})$$

R3.2 Integrals

$$\int f(g(x))g'(x) dx = \int f(y) dy, \quad y = g(x) \text{ and } g'(x) = dy/dx \quad (\text{R3.9})$$

$$\int u dv = uv - \int v du \quad (\text{R3.10})$$

$$\int \frac{f'(x) dx}{f(x)} = \log_e f(x) \quad (\text{R3.11})$$

$$\int \frac{dx}{x} = \log_e x \quad (\text{R3.12})$$

$$\int x^n dx = \frac{x^{n+1}}{n+1} \quad (\text{R3.13})$$

$$\int e^x dx = e^x \quad (\text{R3.14})$$

$$\int a^x dx = \frac{a^x}{\log_e a} \quad (\text{R3.15})$$

$$\int a^{bx} dx = \frac{a^{bx}}{b \log_e a} \quad (\text{R3.16})$$

$$\int \log_e x dx = x \log_e x - x \quad (\text{R3.17})$$

$$\int \sin x dx = -\cos x \quad (\text{R3.18})$$

$$\int \cos x dx = \sin x \quad (\text{R3.19})$$

$$\int \tan x dx = -\log \cos x. \quad (\text{R3.20})$$

R3.3 l'Hôpital's Rule

Consider a fraction $f(x)/g(x)$ for which at $x = x_0$, $f(x_0) = g(x_0) = 0$ (or $f(x_0) = g(x_0) = \infty$). Then

$$\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow x_0} \frac{f'(x)}{g'(x)} \quad (\text{R3.21})$$

as long as the limits on the right-hand side exist and are finite.

R3.4 Examples

Example R3.1. Evaluate the integral

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(\omega) e^{j\omega n} d\omega,$$

where

$$X(\omega) = \begin{cases} \cos(\alpha\omega), & |\omega| \leq \omega_0 \\ 0, & \omega_0 < |\omega| \leq \pi. \end{cases}$$

Answer:

$$\begin{aligned}x[n] &= \frac{1}{2\pi} \int_{-\omega_0}^{\omega_0} \cos(\alpha\omega) e^{j\omega n} d\omega \\&= \frac{1}{2\pi} \int_{-\omega_0}^{\omega_0} \frac{1}{2} (e^{j\alpha\omega} + e^{-j\alpha\omega}) e^{j\omega n} d\omega \\&= \frac{1}{4\pi} \left(\int_{-\omega_0}^{\omega_0} e^{j\alpha\omega} e^{j\omega n} d\omega + \int_{-\omega_0}^{\omega_0} e^{-j\alpha\omega} e^{j\omega n} d\omega \right) \\&= \frac{1}{4\pi} \left(\frac{1}{j(\alpha+n)} e^{j(\alpha+n)\omega} \Big|_{-\omega_0}^{\omega_0} + \frac{1}{j(-\alpha+n)} e^{j(-\alpha+n)\omega} \Big|_{-\omega_0}^{\omega_0} \right) \\&= \frac{1}{4\pi} \left(\frac{1}{j(\alpha+n)} 2j \sin(\alpha+n)\omega_0 + \frac{1}{j(-\alpha+n)} 2j \sin(-\alpha+n)\omega_0 \right) \\&= \frac{\sin(\alpha+n)\omega_0}{2\pi(\alpha+n)} + \frac{\sin(-\alpha+n)\omega_0}{2\pi(-\alpha+n)}.\end{aligned}$$

Example R3.2. Evaluate the integral

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(\omega) e^{j\omega n} d\omega,$$

where

$$X(\omega) = \begin{cases} \omega, & |\omega| \leq \omega_0 \\ 0, & \omega_0 < |\omega| \leq \pi. \end{cases}$$

using integration by parts.

Answer:

$$\begin{aligned}x[n] &= \frac{1}{2\pi} \int_{-\omega_0}^{\omega_0} \omega e^{j\omega n} d\omega \\&= \frac{1}{2\pi} \cdot \frac{1}{jn} \int_{-\omega_0}^{\omega_0} \omega e^{j\omega n} d(jn)\omega \\&= \frac{1}{2\pi} \cdot \frac{1}{jn} \int_{-\omega_0}^{\omega_0} \omega de^{j\omega n} \\&= \frac{1}{2\pi} \cdot \frac{1}{jn} \left[\omega e^{j\omega n} \Big|_{-\omega_0}^{\omega_0} - \int_{-\omega_0}^{\omega_0} e^{j\omega n} d\omega \right] \\&= \frac{1}{2\pi} \cdot \frac{1}{jn} \left[\omega_0 e^{j\omega_0 n} - (-\omega_0) e^{-j\omega_0 n} - \frac{1}{jn} e^{j\omega n} \Big|_{-\omega_0}^{\omega_0} \right] \\&= \frac{1}{2\pi} \cdot \frac{1}{jn} \left[\omega_0 (e^{j\omega_0 n} + e^{-j\omega_0 n}) - \frac{1}{jn} (e^{j\omega_0 n} - e^{-j\omega_0 n}) \right] \\&= \frac{1}{\pi(jn)} \left[\omega_0 \cos(\omega_0 n) - \frac{1}{n} \sin(\omega_0 n) \right].\end{aligned}$$

R4 Complex Numbers

R4.1 Definition

A complex number z is represented in the Cartesian coordinate as

$$z = x + jy$$

where $j = \sqrt{-1}$, and x and y are real numbers and denoted as the real and imaginary parts of z , respectively. The complex numbers can also be represented in polar form as

$$z = |z|e^{j\theta},$$

where $|z|$ and θ are the magnitude and angle of z , respectively:

$$|z| = \sqrt{x^2 + y^2}, \quad \theta = \tan^{-1}\left(\frac{y}{x}\right).$$

The *principal value* of the angle of z is given by

$$-\pi < \theta \leq \pi.$$

Figure R4.1 shows the representation of a complex number z in the complex plane. By using *Euler's Formula* (see Eq.(R1.29)) we can find the representation of a complex number in Cartesian form from its polar form:

$$x = |z| \cos \theta, \quad y = |z| \sin \theta.$$

It should be pointed out here that negative real numbers have angle

$$\theta = (2k + 1)\pi$$

with k any integer.

Example R4.1. Let $z = 2 + j\sqrt{3}$, we can express it in polar form by calculating its magnitude and angle:

$$|z| = \sqrt{2^2 + 3} = \sqrt{7}, \quad \theta = \tan^{-1}\left(\frac{\sqrt{3}}{2}\right).$$

Therefore,

$$z = 2 + j\sqrt{3} = \sqrt{7}e^{j \tan^{-1}\left(\frac{\sqrt{3}}{2}\right)}.$$




Figure R4.1: Representation of a complex number z in Cartesian form and polar form.

The *conjugate* of a complex number in Cartesian form is obtained by negating the imaginary part:

$$z^* = (x + jy)^* = x^* + (jy)^* = x - jy.$$

In polar form, the conjugate is obtained by changing the sign of the angle:

$$z^* = (re^{j\theta})^* = r e^{-j\theta}.$$

R4.2 Complex Arithmetic

(1) Addition and Subtraction

$z_1 = x_1 + jy_1$ and $z_2 = x_2 + jy_2$ be two complex numbers. Then

$$z_1 + z_2 = (x_1 + x_2) + j(y_1 + y_2)$$

where $(x_1 + x_2)$ are $(y_1 + y_2)$ are the real and imaginary parts of the sum $z_1 + z_2$, respectively. Similarly,

$$z_1 - z_2 = (x_1 - x_2) + j(y_1 - y_2)$$

where $(x_1 - x_2)$ are $(y_1 - y_2)$ are the real and imaginary parts of the difference $z_1 - z_2$, respectively.

Example R4.2. Let $z_1 = 1.3 + j5.2$ and $z_2 = 2.7 - j3.6$ then

$$\begin{aligned} z_1 + z_2 &= (1.3 + j5.2) + (2.7 - j3.6) = 4 + j1.6 \\ z_1 - z_2 &= (1.3 + j5.2) - (2.7 - j3.6) = -1.4 + 8.8j. \end{aligned}$$

(2) Multiplication

Let $z_1 = x_1 + jy_1$ and $z_2 = x_2 + jy_2$ then

$$\begin{aligned} z_1 \cdot z_2 &= (x_1 + jy_1)(x_2 + jy_2) \\ &= x_1x_2 + jx_1y_2 + jx_2y_1 + j^2y_1y_2 \\ &= (x_1x_2 - y_1y_2) + j(x_1y_2 + x_2y_1). \end{aligned}$$

Example R4.3. Let $z_1 = 1 + j\sqrt{3}$, $z_2 = 2 - j2$. The product of z_1, z_2 calculated in polar form is given by

$$(1 + j\sqrt{3})(2 - j2) = 2e^{j\pi/3} \cdot 2\sqrt{2}e^{-j\pi/4} = 4\sqrt{2}e^{j\pi/12} = 5.4641 + j1.4641.$$

Calculating in the Cartesian form we get

$$(1 + j\sqrt{3})(2 - j2) = (2 + 2\sqrt{3}) + j(2\sqrt{3} - 2) = 5.4641 + j1.4641.$$

(3) Division

The division of two complex numbers z_0 and z_1 can be carried out either in polar form or in Cartesian form. In the former case

$$w = \frac{z_0}{z_1} = \frac{r_0e^{j\theta_0}}{r_1e^{j\theta_1}} = \frac{r_0}{r_1}e^{j(\theta_0-\theta_1)}.$$

In the latter case

$$w = \frac{z_0}{z_1} = \frac{x_0 + jy_0}{x_1 + jy_1} = \frac{(x_0 + jy_0)(x_1 - jy_1)}{(x_1 + jy_1)(x_1 - jy_1)} = \frac{(x_0x_1 + y_0y_1) + j(x_1y_0 - x_0y_1)}{x_1^2 + y_1^2}.$$

Example R4.4. To divide $2 + j2$ by $1 - j$, we calculate in polar form as follows:

$$\frac{2 + j2}{1 - j} = \frac{2\sqrt{2}e^{j\frac{\pi}{4}}}{\sqrt{2}e^{-j\frac{\pi}{4}}} = 2e^{j(\frac{\pi}{4})-(-\frac{\pi}{4})} = 2e^{j\frac{\pi}{2}} = j2.$$

Calculating in the Cartesian form we get

$$\frac{2 + j2}{1 - j} = \frac{(2 + j2)(1 + j)}{(1 - j)(1 + j)} = \frac{(2 - 2) + j(2 + 2)}{1^2 + 1^2} = j2.$$

(4) Inverse

The inverse of a complex number is a special case of division where the numerator is 1. In polar form we have

$$z^{-1} = \frac{1}{z} = \frac{1}{re^{j\theta}} = \frac{1}{r}e^{-j\theta}.$$

Equivalently, in the Cartesian form we have

$$z^{-1} = \frac{1}{z} = \frac{1}{x + jy} = \frac{x - jy}{(x + jy)(x - jy)} = \frac{x - jy}{x^2 + y^2}.$$

R5 Complex Variables

R5.1 Function of a Complex Variable

A function of the complex variable z can be written as

$$f(z) = u(z) + jv(z)$$

where $u(z)$ and $v(z)$ are real functions of z . In the Cartesian form, we define $z = x + jy$ for real x and y . Therefore, the values of $u(z)$ and $v(z)$ depend on x and y , and we can express the complex function $f(z)$ as

$$f(z) = u(x, y) + jv(x, y).$$

If $z = re^{j\theta}$, then $f(z)$ can be expressed as

$$f(z) = u(r, \theta) + jv(r, \theta),$$

where $u(r, \theta)$ and $v(r, \theta)$ are the real and imaginary parts of $f(z)$.

R5.2 Analytic Function of a Complex Variable

Definition R5.1. A function $f(z)$ is said to be *differentiable* at a point z_0 in the z -plane if the limit

$$f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$$

exists. Note that $f(z_0 + \Delta z)$ can approach $f(z_0)$ along any path. This limit is called the *derivative* of $f(z)$ at point z_0 .

Definition R5.2. A function $f(z)$ of a complex variable z is *analytic* in the region R in the complex z -plane if and only if all the derivatives of $f(z)$ exist at all points inside the region R .

R5.3 Analytic Continuation

If the values of a function $f(z)$ of a complex variable are known everywhere on a closed contour C inside a region R where $f(z)$ is analytic, then the values of $f(z)$ at all points in R can be found by mapping from the contour C to any point in R .

R5.4 Cauchy's Integral Formula

If a function $f(z)$ is analytic both on and inside a counterclockwise closed contour C and if z_0 is any point inside C , then

$$f(z_0) = \frac{1}{2\pi j} \oint_C f(z) \frac{1}{z - z_0} dz, \quad (\text{R5.1})$$

$$f'(z_0) = \frac{1}{2\pi j} \oint_C f(z) \frac{1}{(z - z_0)^2} dz \quad (\text{R5.2})$$

$$f''(z_0) = \frac{2}{2\pi j} \oint_C f(z) \frac{1}{(z - z_0)^3} dz \quad (\text{R5.3})$$

$$\vdots \quad (\text{R5.4})$$

$$f^{(n)}(z_0) = \frac{n!}{2\pi j} \oint_C f(z) \frac{1}{(z - z_0)^{n+1}} dz. \quad (\text{R5.5})$$

where $f'(z_0) = \left. \frac{df(z)}{dz} \right|_{z=z_0}$, $f''(z_0) = \left. \frac{d^2 f(z)}{dz^2} \right|_{z=z_0}$, and $f^{(n)}(z_0) = \left. \frac{d^n f(z)}{dz^n} \right|_{z=z_0}$.

Eq.(R5.1) is often referred as the Cauchy's integral formula.

By combining Eq.(R5.1) - Eq.(R5.5) we arrive at an useful relation:

$$\frac{1}{2\pi j} \oint_C z^{k-1} dz = \begin{cases} 1 & , k = 0 \\ 0 & , k \neq 0 \end{cases} \quad (\text{R5.6})$$

where C is a counterclockwise closed contour encircling $z = 0$.

R5.5 Cauchy's Residue Theorem

If a function $f(z)$ is analytic both on and inside a counterclockwise closed contour C except at poles z_k , $k = 1, 2, \dots, n$, then

$$\frac{1}{2\pi j} \oint_C f(z) dz = \sum_k \left[\text{residue of } f(z) \text{ at pole } z_k \text{ inside } C \right]. \quad (\text{R5.7})$$

In the case when $f(z)$ is a rational function of z and has pole at $z = z_k$ of multiplicity m , we can express $f(z)$ as

$$f(z) = \frac{\Gamma(z)}{(z - z_k)^m},$$

where $\Gamma(z)$ does not have any pole at $z = z_k$. Thus the residue of $f(z)$ at the pole z_k inside C is given by

$$\frac{1}{(m-1)!} \left[\frac{d^{m-1} \Gamma(z)}{dz^{m-1}} \right]_{z=z_k}. \quad (\text{R5.8})$$

R6 Continuous-Time Signals

R6.1 Energy and Power

The *total energy* of a continuous-time signal $x(t)$ is given by

$$E_x = \lim_{T \rightarrow \infty} \int_{-T}^T |x(t)|^2 dt.$$

The *average power* of a continuous-time $x(t)$ is given by

$$P_x = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T |x(t)|^2 dt.$$

The definition of total energy can be explained as the area under the squared signal $|x(t)|^2$, and it is a measurement of the strength of the signal $x(t)$ over infinite time. However, there are signals with infinite energy so we need to evaluate the average power of the signal $x(t)$ as a measurement of the strength over one unit time.

R6.2 Continuous-Time Sinusoidal and Exponential Signals

R6.2.1 Definition

The continuous-time *real sinusoidal signal* with constant amplitude is of the form

$$x(t) = A \cos(\Omega_0 t + \phi), \quad (\text{R6.1})$$

where A, Ω_0 and ϕ are real numbers. The parameters A, Ω_0 and ϕ are called, respectively, the *amplitude*, the *angular frequency*, and the *phase* of the sinusoidal signal $x(t)$.

The *complex exponential signal* is expressed in the form

$$x(t) = A\alpha^t, \quad (\text{R6.2})$$

where

$$\alpha = e^{\sigma_0 + j\Omega_0}, A = |A| e^{j\phi}. \quad (\text{R6.3})$$

If A and α are both real, the signal of Eq. (R6.2) reduces to *real exponential signal*. For $t \geq 0$ such a signal with $|\alpha| < 1$ decays exponentially as t increases and with $|\alpha| > 1$ grows exponentially as t increases.

In addition, we can rewrite Eq. (R6.2) as

$$x(t) = Ae^{(\sigma_0 + j\Omega_0)t} = |A| e^{\sigma_0 t} e^{j(\Omega_0 t + \phi)} \quad (\text{R6.4})$$

$$= |A| e^{\sigma_0 t} \cos(\Omega_0 t + \phi) + j|A| e^{\sigma_0 t} \sin(\Omega_0 t + \phi). \quad (\text{R6.5})$$

Thus the real and imaginary parts of a complex exponential signal are real sinusoidal signals.

The *fundamental period* T_0 of a complex exponential signal (Eq. (R6.4)) with $\sigma_0 = 0$ is defined to be the smallest positive T_0 satisfying

$$|A| e^{j(\Omega_0 t + \phi)} = |A| e^{j(\Omega_0(t+T_0) + \phi)}, \quad (\text{R6.6})$$

or equivalently,

$$e^{j\Omega_0 T_0} = 1. \quad (\text{R6.7})$$

Therefore,

$$T_0 = \frac{2\pi}{|\Omega_0|}. \quad (\text{R6.8})$$

R6.2.2 Properties

The properties of continuous-time sinusoidal and exponential signals and comparisons with discrete-time sinusoidal and exponential sequences are discussed as follows.

1. Periodicity for any choice of Ω_0

Note that the continuous-time sinusoidal signal $A \cos(\Omega_0 t + \phi)$ (Eq. (R6.1)) and the continuous-time complex exponential signal $|A| e^{j(\Omega_0 t + \phi)}$ are periodic signals of any choice of Ω_0 . However, discrete-time sequences are not always periodic with any choice of ω_0 . Discrete sinusoidal sequence $A \cos(\omega_0 n + \phi)$ and discrete complex exponential sequence $|A| e^{j(\omega_0 n + \phi)}$ are periodic with period N only if $\omega_0 N$ is an integer multiple of 2π , i.e., $\omega_0 N = 2\pi r$ where N and r are positive integers. For example, $\cos(\frac{\pi n}{4})$ is a periodic sequence while $\cos(\frac{n}{4})$ is not periodic.

2. Distinctness for different Ω_0, Ω_1

Any two continuous-time sinusoidal signals

$$A \cos(\Omega_0 t + \phi), A \cos(\Omega_1 t + \phi), \Omega_0 \neq \Omega_1$$

have different waveforms. Similarly, any two continuous-time exponential signals with $\Omega_0 \neq \Omega_1$ also have different waveforms. Unlike the continuous-time case, discrete-time sinusoidal sequences

$$A \cos(\omega_0 n + \phi), A \cos(\omega_1 n + \phi), \omega_0 = \omega_1 + 2\pi k$$

have the same sequence values. Similarly, any two discrete-time exponential sequences with $\omega_0 = \omega_1 + 2\pi k$ also have the same sequence values.

R6.3 Continuous-Time Eigenfunction

If the input signal of any LTI system has output signal being the input signal multiplied by a complex constant, this certain type of input signal is called the *eigenfunction* and the complex constant is called the *eigenvalue*.

Example R6.1. We want to show that the complex exponential signal defined in Eq. (R6.2)

$$x(t) = A\alpha^t$$

is an eigenfunction of an LTI continuous-time system with an impulse response $h(t)$.

By using the convolution integral, we have

$$\begin{aligned} y(t) &= \int_{-\infty}^{\infty} h(\tau) A \alpha^{(t-\tau)} d\tau \\ &= \left(\int_{-\infty}^{\infty} h(\tau) \alpha^{-\tau} d\tau \right) A \alpha^t. \end{aligned}$$

Since the integral inside the brackets is independent of t , we can therefore say that the input signal $A\alpha^t$ is an eigenfunction.

Example R6.2. We want to show that the sum of any two complex exponential signals

$$x(t) = A\alpha^t + B\beta^t$$

is not an eigenfunction of an LTI continuous-time system with an impulse response $h(t)$.

By using the convolution integral, we have

$$\begin{aligned} y(t) &= \int_{-\infty}^{\infty} h(\tau) \left(A\alpha^{(t-\tau)} + B\beta^{(t-\tau)} \right) d\tau \\ &= \left(\int_{-\infty}^{\infty} h(\tau) \alpha^{-\tau} d\tau \right) A\alpha^t + \left(\int_{-\infty}^{\infty} h(\tau) \beta^{-\tau} d\tau \right) B\beta^t. \end{aligned}$$

Since the input signal $x(t)$ cannot be extracted from the summation above, we can therefore say that the input signal $A\alpha^t + B\beta^t$ is not an eigenfunction.

R6.4 Continuous-Time Fourier Series

R6.4.1 Definition

Given a periodic continuous-time signal $x(t)$ with period T_0 and fundamental frequency $\Omega_0 = 2\pi/T_0$, the Fourier series expansion of $x(t)$ is given by the linear combination of the set of harmonically related complex exponentials

$$e^{jk\Omega_0 t} = e^{jk\frac{2\pi}{T_0} t}, \quad k = 0, \pm 1, \pm 2, \dots,$$

i.e.,

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{jk\Omega_0 t} = \sum_{k=-\infty}^{\infty} a_k e^{jk\frac{2\pi}{T_0} t} \quad (\text{R6.9})$$

$$a_k = \frac{1}{T_0} \int_{T_0} x(t) e^{-jk\Omega_0 t} dt = \frac{1}{T_0} \int_{T_0} x(t) e^{-jk\frac{2\pi}{T_0} t} dt. \quad (\text{R6.10})$$

Note that the notation \int_{T_0} denotes the integration over any interval of length T_0 . The Eq.(R6.9) is referred to as the *synthesis* equation and the Eq.(R6.10) is referred to as the *analysis* equation. The coefficient a_k is called the *Fourier series coefficient*.

Table R6.1: Properties of Continuous-Time Fourier Series.

Type of Property	Periodic Signal with frequency $\Omega_0 = 2\pi/T$	Fourier Series Coefficients
	$g(t)$	a_k
	$h(t)$	b_k
Linearity	$\alpha g(t) + \beta h(t)$	$\alpha a_k + \beta b_k$
Time Shifting	$g(t - t_0)$	$a_k e^{-jk\Omega_0 t_0}$
Frequency Shifting	$e^{jM\Omega_0 t} g(t)$	a_{k-M}
Multiplication	$g(t)h(t)$	$\sum_{l=-\infty}^{\infty} a_l b_{k-l}$
Time Reversal	$g(-t)$	a_{-k}
Conjugation	$g^*(t)$	a_{-k}^*
Time Scaling	$g(\alpha t), \alpha > 0$	a_k
Periodic Convolution	$\int_T g(\tau)h(t - \tau) d\tau$	$T a_k b_k$

Example R6.3. Find the Fourier series coefficients of the continuous-time signal

$$x(t) = 1 + \cos(\Omega_0 t) + 2 \cos(2\Omega_0 t + \frac{\pi}{3}) + 4 \sin(3\Omega_0 t - \frac{\pi}{4})$$

with fundamental frequency Ω_0 .

By using the Euler's Formula, it can be shown that

$$\begin{aligned} x(t) &= 1 + \frac{1}{2}(e^{j\Omega_0 t} + e^{-j\Omega_0 t}) + \frac{2}{2}(e^{j(2\Omega_0 t + \frac{\pi}{3})} + e^{-j(2\Omega_0 t + \frac{\pi}{3})}) + \frac{4}{2j}(e^{j(3\Omega_0 t - \frac{\pi}{4})} - e^{-j(3\Omega_0 t - \frac{\pi}{4})}) \\ &= 1 + \frac{1}{2}e^{j\Omega_0 t} + \frac{1}{2}e^{-j\Omega_0 t} + e^{j\frac{\pi}{3}}e^{j2\Omega_0 t} + e^{-j\frac{\pi}{3}}e^{-j2\Omega_0 t} + \frac{2}{j}e^{-j\frac{\pi}{4}}e^{j3\Omega_0 t} - \frac{2}{j}e^{j\frac{\pi}{4}}e^{-j3\Omega_0 t}. \end{aligned}$$

Therefore, the Fourier series coefficients are

$$\begin{aligned}
a_0 &= 1, \\
a_1 &= \frac{1}{2}, \quad a_{-1} = \frac{1}{2}, \\
a_2 &= e^{j\frac{\pi}{3}} = \frac{1 + j\sqrt{3}}{2}, \quad a_{-2} = e^{-j\frac{\pi}{3}} = \frac{1 - j\sqrt{3}}{2}, \\
a_3 &= \frac{2}{j}e^{-j\frac{\pi}{4}} = \sqrt{2}(-1 - j), \quad a_{-3} = -\frac{2}{j}e^{j\frac{\pi}{4}} = \sqrt{2}(-1 + j), \\
a_k &= 0, |k| > 3.
\end{aligned}$$

Example R6.4. Find the Fourier series coefficients of the impulse train

$$x(t) = \sum_{k=-\infty}^{\infty} \delta(t - kT_0)$$

with period T_0 .

By calculating Eq.(R6.10) in the interval $-T_0/2 \leq t \leq T_0/2$, we can get

$$a_k = \frac{1}{T_0} \int_{-T_0/2}^{T_0/2} \delta(t) e^{-jk\frac{2\pi}{T_0}t} dt = \frac{1}{T_0}.$$

Therefore, all the Fourier series coefficients of the impulse train have the same value $1/T_0$.

Some important properties of continuous-time Fourier series are listed in Table R6.1 for quick reference.

R6.4.2 Dirichlet Conditions

In order to verify the existence of Fourier series representation for a periodic signal $x(t)$, we need to examine the Dirichlet conditions. The Dirichlet conditions are given by:

1. $x(t)$ must be absolutely integrable, i.e.,

$$\int_{T_0} |x(t)| dt < \infty$$

2. In any finite interval of time, $x(t)$ has a finite number of local maxima and local minima.
3. In any finite interval of time, $x(t)$ has a finite number of discontinuities.

The Dirichlet conditions guarantee that $x(t)$ equals its Fourier series representation

$$\sum_{k=-\infty}^{\infty} a_k e^{jk\Omega_0 t}$$

at all values of t except at discontinuities of $x(t)$. Note that Dirichlet conditions are only sufficient but not necessary conditions.

R6.5 Continuous-Time Fourier Transform

Given an aperiodic continuous-time signal $x(t)$, the continuous-time Fourier transform of $x(t)$ is given by

$$X(j\Omega) = \int_{-\infty}^{\infty} x(t) e^{-j\Omega t} dt \quad (\text{R6.11})$$

$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\Omega) e^{j\Omega t} d\Omega. \quad (\text{R6.12})$$

The transform $X(j\Omega)$ is referred to as the *spectrum* of $x(t)$ because it provides the information of $x(t)$ when evaluated by complex exponential signals at different frequencies.

Some important properties of continuous-time Fourier transform are listed in Table [R6.2](#) for quick reference.

Table R6.2: Properties of Continuous-Time Fourier Transform.

Property	Signal	Fourier Transform
	$g(t)$	$G(j\Omega)$
	$h(t)$	$H(j\Omega)$
Linearity	$\alpha g(t) + \beta h(t)$	$\alpha G(j\Omega) + \beta H(j\Omega)$
Time Shifting	$g(t - t_0)$	$G(j\Omega)e^{-j\Omega t_0}$
Frequency Shifting	$e^{j\Omega_0 t} g(t)$	$G(j(\Omega - \Omega_0))$
Multiplication	$g(t)h(t)$	$\frac{1}{2\pi}G(j\Omega) * H(j\Omega)$
Time Reversal	$g(-t)$	$G(-j\Omega)$
Conjugation	$g^*(t)$	$G^*(-j\Omega)$
Time Scaling	$g(\alpha t)$	$\frac{1}{ \alpha }G\left(\frac{j\Omega}{\alpha}\right)$
Convolution	$g(t) * h(t)$	$G(j\Omega)H(j\Omega)$
Differentiation in Time	$\frac{d}{dt}g(t)$	$j\Omega G(j\Omega)$
Integration	$\int_{-\infty}^t g(\tau) d\tau$	$\frac{1}{j\Omega}G(j\Omega) + \pi G(0)\delta(\Omega)$
Real and Even in Time	$g(t)$ real and even	$G(j\Omega)$ real and even
Real and Odd in Time	$g(t)$ real and odd	$G(j\Omega)$ purely imaginary and odd

R7 Discrete Fourier Series

Given a periodic sequence $x[n]$ with period N , the fundamental period is defined to be the smallest integer N such that $x[n] = x[n + N]$ is satisfied, and the fundamental frequency is defined to be $\omega_0 = 2\pi/N$. The harmonics are sequences whose frequencies are integer multiples of the fundamental frequency. For discrete complex exponential signals, the k -th harmonic is expressed as

$$e^{jk\omega_0 n} = e^{jk\frac{2\pi}{N}n}, \quad k = 0, \pm 1, \pm 2, \dots$$

Note that there are only N distinct harmonics for discrete complex exponential signals with fundamental frequency $\omega_0 = 2\pi/N$ because every two signals with frequencies which differ in $2\pi m$ have the same waveform, i.e.,

$$e^{jk(\omega_0 + 2m\pi)n} = e^{jk(\frac{2\pi}{N} + 2m\pi)n} = e^{jk(\frac{2\pi}{N})n} \cdot e^{jk2mn\pi} = e^{jk(\frac{2\pi}{N})n}.$$

The discrete Fourier series expansion of periodic signal $x[n]$ is the expression in form of a linearly weighted combination of a fundamental and a series of harmonic complex exponential signals.

$$x[n] = \sum_{k=0}^{N-1} a_k e^{jk\omega_0 n} = \sum_{k=0}^{N-1} a_k e^{jk\frac{2\pi}{N}n},$$

where

$$a_k = \frac{1}{N} \sum_{n=0}^{N-1} x[n] e^{-jk\omega_0 n} = \frac{1}{N} \sum_{n=0}^{N-1} x[n] e^{-jk\frac{2\pi}{N}n}.$$

Example R7.1. Calculate the Fourier series coefficients a_k for the following periodic signal

$$\{x[n]\} = \{\dots, 1, 1, 1, 1, 0, 0, \dots\}.$$

↑

We can observe that $N = 6$ so

$$\begin{aligned} a_k &= \frac{1}{6} \sum_{n=0}^5 x[n] e^{-jk(\frac{2\pi}{6})n} \\ &= \frac{1}{6} (e^{-jk(\frac{2\pi}{6})0} + e^{-jk(\frac{2\pi}{6})1} + e^{-jk(\frac{2\pi}{6})2} + e^{-jk(\frac{2\pi}{6})3} + 0 + 0) \\ &= \frac{1}{6} (1 + e^{-jk\frac{\pi}{3}} + e^{-jk\frac{2\pi}{3}} + e^{-jk\pi}) \\ &= \frac{1}{6} (1 + e^{-jk\frac{\pi}{3}} + (-1)^k e^{jk\frac{\pi}{3}} + (-1)^k). \end{aligned}$$

Example R7.2. Calculate signal $x[n]$ from the following Fourier series coefficients a_k

$$\{a_k\} = \{\dots, 1/4, 1/2, 1, 1/2, 1/4, 0, 1/4, 1/2, 1, 1/2, \dots\}.$$

↑ .

We can observe that $N = 6$ so

$$\begin{aligned} x[n] &= \sum_{k=\langle N \rangle} a_k e^{jk\frac{2\pi}{6}n} = 1 + \frac{1}{2} e^{j\frac{2\pi}{6}n} + \frac{1}{4} e^{j\frac{2\pi}{6}2n} + 0 + \frac{1}{4} e^{j\frac{2\pi}{6}4n} + \frac{1}{2} e^{j\frac{2\pi}{6}5n} \\ &= 1 + \frac{1}{2} e^{j\frac{\pi n}{3}} + \frac{1}{4} e^{j\frac{2\pi n}{3}} + 0 + \frac{1}{4} e^{j\frac{4\pi n}{3}} + \frac{1}{2} e^{j\frac{5\pi n}{3}} \\ &= 1 + \frac{1}{2} e^{j\frac{\pi n}{3}} + \frac{1}{4} e^{j\frac{2\pi n}{3}} + 0 + \frac{1}{4} e^{j(2\pi n - \frac{2\pi n}{3})} + \frac{1}{2} e^{j(2\pi n - \frac{\pi n}{3})} \\ &= 1 + \frac{1}{2} e^{j\frac{\pi n}{3}} + \frac{1}{4} e^{j\frac{2\pi n}{3}} + 0 + \frac{1}{4} e^{-j\frac{2\pi n}{3}} + \frac{1}{2} e^{-j\frac{\pi n}{3}} \\ &= 1 + \cos\left(\frac{\pi n}{3}\right) + \frac{1}{2} \cos\left(\frac{2\pi n}{3}\right). \end{aligned}$$

R8 Matrix Algebra

R8.1 Definition

A matrix is a rectangular array of real or complex numbers enclosed in brackets; for instance,

$$\begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 5 & 6 \\ 2 & 1 & -3 \end{bmatrix}, \begin{bmatrix} 3j & 5 \\ 2-4j & 1+j \\ 7 & 5-3j \end{bmatrix}.$$

A matrix with K rows and M columns is called a $K \times M$ matrix. For example, the matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

is a 3×3 matrix. The matrix

$$\begin{bmatrix} 1 & 4 \\ 2 & 6 \\ -3j & 1 \\ 1+j & 1 \end{bmatrix}$$

is a 4×2 matrix. The matrix

$$\mathbf{U} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1M} \\ a_{21} & a_{22} & \cdots & a_{2M} \\ \vdots & \vdots & \ddots & \vdots \\ a_{K1} & a_{K2} & \cdots & a_{KM} \end{bmatrix} \quad (\text{R8.1})$$

is a $K \times M$ matrix and the number a_{rs} , $r = 1, 2, \dots, K$, and $s = 1, 2, \dots, M$ is called the entry of \mathbf{U} .

R8.2 Transpose

The transpose, \mathbf{U}^T , of a $K \times M$ matrix \mathbf{U} is the $M \times K$ matrix formed by interchanging the rows and columns of \mathbf{U} . For example, the transpose of the

matrix \mathbf{U} given in Eq.(R8.1) is a $M \times K$ matrix given by

$$\mathbf{U}^T = \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{K1} \\ a_{12} & a_{22} & \cdots & a_{K2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1M} & a_{2M} & \cdots & a_{KM} \end{bmatrix}. \quad (\text{R8.2})$$

R8.3 Toeplitz Matrix

The $N \times N$ matrix \mathbf{U} is a Toeplitz matrix if all entries along the line parallel to the main diagonal are the same. For example,

$$\mathbf{U} = \begin{bmatrix} a_0 & a_{-1} & a_{-2} & a_{-3} \\ a_1 & a_0 & a_{-1} & a_{-2} \\ a_2 & a_1 & a_0 & a_{-1} \\ a_3 & a_2 & a_1 & a_0 \end{bmatrix}$$

is a 4×4 Toeplitz matrix.

R8.4 Circulant Matrix

The $N \times N$ matrix \mathbf{U} is a circulant matrix if each row equals the right circular shift of the previous row by one entry. For example,

$$\mathbf{U} = \begin{bmatrix} a_0 & a_1 & a_2 & a_3 \\ a_3 & a_0 & a_1 & a_2 \\ a_2 & a_3 & a_0 & a_1 \\ a_1 & a_2 & a_3 & a_0 \end{bmatrix}$$

is a 4×4 Circulant matrix.

R8.5 Determinant

If the 2×2 matrix \mathbf{U} is

$$\mathbf{U} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}, \quad (\text{R8.3})$$

then the determinant of the \mathbf{U} is given by

$$\det(\mathbf{U}) = a_{11}a_{22} - a_{12}a_{21}. \quad (\text{R8.4})$$

Example R8.1. The determinant of the matrix

$$\mathbf{U} = \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}$$

is $\det(\mathbf{U}) = 3 \cdot 2 - 4 \cdot 1 = 6 - 4 = 2$.

If the 3×3 matrix \mathbf{U} is

$$\mathbf{U} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}, \quad (\text{R8.5})$$

then the determinant of \mathbf{U} is given by

$$\det(\mathbf{U}) = a_{11}a_{22}a_{33} + a_{21}a_{32}a_{13} + a_{12}a_{23}a_{31} - a_{13}a_{22}a_{31} - a_{11}a_{32}a_{23} - a_{12}a_{21}a_{33}. \quad (\text{R8.6})$$

Example R8.2. The determinant of the matrix

$$\begin{bmatrix} 1 & 4 & -6 \\ 2 & 1 & 3 \\ 4 & 5 & -2 \end{bmatrix}$$

is

$$\begin{aligned} \det(\mathbf{U}) &= 1 \cdot 1 \cdot (-2) + 2 \cdot 5 \cdot (-6) + 4 \cdot 3 \cdot 4 - (-6) \cdot 1 \cdot 4 - 1 \cdot 5 \cdot 3 - 4 \cdot 2 \cdot (-2) \\ &= (-2) + (-60) + 48 - (-24) - 15 - (-16) = 11. \end{aligned}$$

If the $N \times N$ matrix \mathbf{U} is

$$\mathbf{U} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1N} \\ a_{21} & a_{22} & \cdots & a_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ a_{N1} & a_{N2} & \cdots & a_{NN} \end{bmatrix}, \quad (\text{R8.7})$$

then the determinant of \mathbf{U} is

$$\det(\mathbf{U}) = a_{r1}(-1)^{r+1}M_{r1} + a_{r2}(-1)^{r+2}M_{r2} + \cdots + a_{rN}(-1)^{r+N}M_{rN} \quad (\text{R8.8})$$

$$= a_{1s}(-1)^{1+s}M_{1s} + a_{2s}(-1)^{2+s}M_{2s} + \cdots + a_{Ns}(-1)^{N+s}M_{Ns} \quad (\text{R8.9})$$

where $r, s = 1$ or 2 or $3 \cdots$, or N and M_{rs} is the minor of a_{rs} (see Section [R8.6](#)).

Example R8.3. To calculate the determinant of the matrix

$$\mathbf{U} = \begin{bmatrix} 1 & 6 & -4 & 6 \\ -7 & 1 & 3 & -6 \\ 2 & -3 & 6 & 5 \\ -2 & 4 & 2 & 6 \end{bmatrix}, \quad (\text{R8.10})$$

we first calculate the minors

$$\begin{aligned} M_{11} &= \begin{bmatrix} 1 & 3 & -6 \\ -3 & 6 & 5 \\ 4 & 2 & 6 \end{bmatrix} = 320, & M_{12} &= \begin{bmatrix} -7 & 3 & -6 \\ 2 & 6 & 5 \\ -2 & 2 & 6 \end{bmatrix} = -344 \\ M_{13} &= \begin{bmatrix} -7 & 1 & -6 \\ 2 & -3 & 5 \\ -2 & 4 & 6 \end{bmatrix} = 232, & M_{14} &= \begin{bmatrix} -7 & 1 & 3 \\ 2 & -3 & 6 \\ -2 & 4 & 2 \end{bmatrix} = 200. \end{aligned}$$

The determinant is therefore given by

$$\begin{aligned} \det(\mathbf{U}) &= 1 \cdot (-1)^{1+1} \cdot 320 + 6 \cdot (-1)^{1+2} \cdot (-344) + (-4) \cdot (-1)^{1+3} \cdot 232 + 6 \cdot (-1)^{1+4} \cdot 200 \\ &= 320 + 2064 - 928 - 1200 = 256. \end{aligned}$$

R8.6 Minor and Cofactor

From \mathbf{U} given in Eq.(R8.7), the minor M_{rs} of a_{rs} in \mathbf{U} is defined to be the determinant of the $(N-1) \times (N-1)$ matrix formed by deleting the r -th row and s -th column of \mathbf{U} . For example,

$$M_{11} = \begin{vmatrix} a_{22} & a_{23} & \cdots & a_{2N} \\ a_{32} & a_{33} & \cdots & a_{3N} \\ \vdots & \vdots & \ddots & \vdots \\ a_{N2} & a_{N3} & \cdots & a_{NN} \end{vmatrix}, \quad M_{12} = \begin{vmatrix} a_{21} & a_{23} & \cdots & a_{2N} \\ a_{31} & a_{33} & \cdots & a_{3N} \\ \vdots & \vdots & \ddots & \vdots \\ a_{N1} & a_{N3} & \cdots & a_{NN} \end{vmatrix}.$$

The cofactor C_{rs} of a_{rs} in \mathbf{U} given in Eq.(R8.7) is defined to be

$$C_{rs} = (-1)^{r+s} M_{rs}. \quad (\text{R8.11})$$

R8.7 Inverse of a Matrix

By Eq.(R8.7), if $\det(\mathbf{U}) \neq 0$, then the inverse of \mathbf{U} exists and is uniquely given by

$$\mathbf{U}^{-1} = \frac{1}{\det(\mathbf{U})} \begin{bmatrix} C_{11} & C_{21} & \cdots & C_{N1} \\ C_{12} & C_{22} & \cdots & C_{N2} \\ \vdots & \vdots & \ddots & \vdots \\ C_{1N} & C_{2N} & \cdots & C_{NN} \end{bmatrix}, \quad (\text{R8.12})$$

where $C_{rs} = (-1)^{r+s}M_{rs}$ is the cofactor of a_{rs} in \mathbf{U} given in Eq.(R8.7).

Example R8.4. In this example we want to find the inverse of the matrix given in Eq.(R8.10). The cofactors are calculated as follows:

$$\begin{aligned} C_{11} &= (-1)^{1+1}M_{11} = 320, & C_{12} &= (-1)^{1+2}M_{12} = 344, \\ C_{13} &= (-1)^{1+3}M_{13} = 232, & C_{14} &= (-1)^{1+4}M_{14} = -200 \\ C_{21} &= (-1)^{2+1}M_{21} = 176, & C_{22} &= (-1)^{2+2}M_{22} = 210, \\ C_{23} &= (-1)^{2+3}M_{23} = 158, & C_{24} &= (-1)^{2+4}M_{24} = -134 \\ C_{31} &= (-1)^{3+1}M_{31} = 240, & C_{32} &= (-1)^{3+2}M_{32} = 234, \\ C_{33} &= (-1)^{3+3}M_{33} = 198, & C_{34} &= (-1)^{3+4}M_{34} = -142 \\ C_{41} &= (-1)^{4+1}M_{41} = -344, & C_{42} &= (-1)^{4+2}M_{42} = -329, \\ C_{43} &= (-1)^{4+3}M_{43} = -239, & C_{44} &= (-1)^{4+4}M_{44} = 227. \end{aligned}$$

Therefore, the inverse is

$$\mathbf{U}^{-1} = \frac{1}{256} \begin{bmatrix} 320 & 176 & 240 & -344 \\ 344 & 210 & 234 & -329 \\ 232 & 158 & 198 & -239 \\ -200 & -134 & -142 & 227 \end{bmatrix}.$$

R8.8 Unitary Matrix and Orthogonal Matrix

The $N \times N$ matrix \mathbf{U} is said to be unitary if

$$\mathbf{U}^H \mathbf{U} = \mathbf{U} \mathbf{U}^H = k\mathbf{I}, \quad (\text{R8.13})$$

where k is any nonzero constant and $\mathbf{U}^H = (\mathbf{U}^T)^*$ is the conjugate-transpose of \mathbf{U} . Note that the unitary matrix is always invertible and $\mathbf{U}^H = \mathbf{U}^{-1}$.

A real unitary matrix \mathbf{U} is also called an orthogonal matrix, i.e.,

$$\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = kI, \quad (\text{R8.14})$$

where k is any nonzero constant and \mathbf{U}^T is the transpose of \mathbf{U} . Similarly, the orthogonal matrix is always invertible and $\mathbf{U}^H = \mathbf{U}^{-1}$. If $k = 1$, then the matrix \mathbf{U} is said to be orthonormal.

R8.9 Cramer's Rule

Consider the set of N linear equations

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1N}x_N &= b_1, \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2N}x_N &= b_2, \\ &\vdots \\ a_{N1}x_1 + a_{N2}x_2 + \cdots + a_{NN}x_N &= b_N, \end{aligned} \quad (\text{R8.15})$$

writing in matrix form yields

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1N} \\ a_{21} & a_{22} & \cdots & a_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ a_{N1} & a_{N2} & \cdots & a_{NN} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_N \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_N \end{bmatrix}. \quad (\text{R8.16})$$

Let D be the determinant of the coefficient matrix

$$D = \begin{vmatrix} a_{11} & a_{12} & \cdots & a_{1N} \\ a_{21} & a_{22} & \cdots & a_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ a_{N1} & a_{N2} & \cdots & a_{NN} \end{vmatrix}. \quad (\text{R8.17})$$

If $D \neq 0$, then the system (R8.15) has the unique solution

$$x_1 = \frac{D_1}{D}, \quad x_2 = \frac{D_2}{D}, \quad \cdots, \quad x_N = \frac{D_N}{D}, \quad (\text{R8.18})$$

where

$$D_1 = \begin{vmatrix} b_1 & a_{12} & \cdots & a_{1N} \\ b_2 & a_{22} & \cdots & a_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ b_N & a_{N2} & \cdots & a_{NN} \end{vmatrix}, \quad D_2 = \begin{vmatrix} a_{11} & b_1 & \cdots & a_{1N} \\ a_{21} & b_2 & \cdots & a_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ a_{N1} & b_N & \cdots & a_{NN} \end{vmatrix}, \quad \cdots, \quad \text{etc.}$$