March 2015

Q.P. Code: 21948

		v	(3 Hours)	Total Marks: 70
ľ	N.B.: (1) (2) (3)	Figures to the	are compulsary e right indicate full marks elled diagram wherever necessary	
1	(a) Answe	er the following	(any seven)	, ,
1.	(i)		length of radiation	5
	(ii)	What is an au		
	(iii)	Name any tw	o types of atomization techniques in	AAS .
	(iv)	Name any on	e non proportional counter for radio	activity measurement
	(v)	Name one rea	agent used for converting non fluore	escent compound to
		fluorescent d		
	(vi)	Give formula	a for determination of correlation co	pefficient
	(vii)	Enlist variou	s bending vibrations in IR spectrosco	opy -
		Define the ur	nit curie used in radiochemistry	
		er the following	(any four)	
	(i)	Discuss mate	erial used for preparation of sample of	ell in UV, visible and
		IR spectrosco	opic analysis	
	(ii)	Compare UV	spectvoscopy and fluorimetry with	respect to sensitivity
		and selectivit	ty .	
	(iii)	What are cat	ionic interferences in AES ? Give r	nethods to overcome
		them		
	(iv)		elength maximum (λ max)? How it	can be determined by
			rophotometer?	
	(v)	Explain the to	erm 'F test'	
			. Ø= .v	0
2.		r the following	(any two)	O
	(i)		nt types of detectors used in IR spect	roscopy. Explain any
	* . * . *	one in detail		141 6
	(ii)	Discuss Stok	es, antistokes and Rayleigh scatteri	ng with reference to
		Raman spectr	roscopy. Support your answer with en	nergy level diagram.
	(iii)	Discuss mea	asurement of equilibrium constant	it using ov visible

(b) What is radionuclidic impurity? Give any one example of the same. Name any

[TURN OVER

spectroscopy

one technique used to determine the same

(b) What are X - rays? What is X - ray diffraction? What are Miller indices?

2 applications of Raman Spectroscopy

- (a) Answer the following (any two)(i) Explain various techniques for handling solid samples in IR spectroscopy.
 - (ii) Draw a neat labelled block diagram of a filter fluorimeter. Expain the function of filters in it.
- (iii) Write principle involved in AAS. Write any two applications of the same
 (b) Calculate concentration of drug A in μg/ml. if it gives absorbance of 0.712 in a 3 4 cm cell, at λmax 254 nm. [Given Molar absorptivity at 254nm = 5229,

Molecular weight of drug A = 248]

5.