139

-1-

**QP Code: 31655** 

## (3 hours)

Max. Marks: 80	20.00
1. Question no.1 is compulsory	رى ن
2. Solve any 3 questions out of remaining	,6
3. Assume data wherever necessary and clearly mention the assumptions made.	20,
4. Draw neat figures as required.	25
Solve any four	20
Define specific force and explain specific force curve in detail.	
Explain S1, S2, S3 profile in detail with neat sketch.	
What are the defects in Lacey's theory?	
Differentiate between streamlined body and bluff body.	
on the verge of separation.	
Assuming that the velocity distribution in the boundary layer is given by $\frac{1}{1}$	10
$\frac{1}{V} = \left(\frac{\delta}{\delta}\right)$ calculate $\frac{\delta}{\delta}, \frac{\delta}{\delta}$ and $\frac{\delta E}{\delta}$ . If at a certain section, free stream velocity V was	
observed to be 10 m/s and the thickness of the boundary layer as 25 mm, then calculate	
$\sim$	
Derive Von Karman's momentum integral Equation.	10
A cylinder 1.2 m in diameter is rotated about its axis in air having a velocity of 128 km	10
	10
The density of the metallic ball is 12000 kg/m. Find:	
(i) The drag force exerted by fluid on metallic ball.	
(ii) The pressure drag and skin friction drag.	
F)	
(5)	P.T.O
	1. Question no.1 is compulsory  2. Solve any 3 questions out of remaining  3. Assume data wherever necessary and clearly mention the assumptions made.  4. Draw neat figures as required.  Solve any four  Define specific force and explain specific force curve in detail.  Explain S1, S2, S3 profile in detail with neat sketch.  What are the defects in Lacey's theory?  Differentiate between streamlined body and bluff body.  How will you determine whether a boundary layer flow is attached flow, detached flow or on the verge of separation.  Assuming that the velocity distribution in the boundary layer is given by $\frac{v}{v} = \left(\frac{v}{\delta}\right)^{1/7} \text{ calculate } \frac{\delta^*}{\delta} \cdot \frac{\theta}{\delta} \text{ and } \frac{\delta E}{\delta}. \text{ If at a certain section, free stream velocity V was observed to be 10 m/s and the thickness of the boundary layer as 25 mm, then calculate the energy loss per unit length due to the formation of boundary layer. Take \rho = 1.226 \text{ kg/m}^3  Derive Von Karman's momentum integral Equation.  A cylinder 1.2 m in diameter is totated about its axis in air having a velocity of 128 km per hour. A lift of 5886 N per meter length of the cylinder is developed on the body. Assuming ideal fluid theory, find rotational speed and the location of the stagnation points. Take \rho = 1.236 \text{ kg/m}^3  A metallic ball of diameter 2 X 10-3 drops in a fluid of sp.gr. 0.95 and viscosity 15 poise. The density of the metallic ball is 12000 kg/m³. Find:$

FW-Con.10926-16